

The Air Plant Guide

Tillandsia species



How to use this guide

The feel of the leaves. The speed of drying. The color at the base. The amount of silver on the surface. Good care begins by noticing which air plant you have and how its room behaves.

THE ONE-CHANGE RULE

Change one condition.
Record what happened. Give
the plant time to answer.

Contents

PLANT IDENTITY	
BOTANICAL	<i>Tillandsia</i> species
FAMILY	Bromeliaceae
ORIGIN	Tropical and subtropical Americas
HABIT	Mostly epiphytic rosette-forming herbs

SECTION	PAGE
Plant profile	04
Care at a glance	05
Light	06
Placement	07
Watering	08
Humidity and temperature	09
Display and feeding	10
Air plant anatomy	11

SECTION	PAGE
Drying and grooming	12
Troubleshooting	13
Mounting and remounting	14
Propagation	15
Seasonal care	16
Pet safety	17
Observation log	18

HOW TO USE THIS GUIDE

Start with the condition you can observe. Avoid making several changes at once.
The printable log at the end is part of the care system, not an afterthought.



Plant profile

Tillandsia is a large American bromeliad genus. Most cultivated air plants are epiphytes: roots anchor them to a support while specialized leaf trichomes absorb water and nutrients. Silver, fuzzy xeric species and greener mesic species do not dry at the same rate.

Care at a glance

Bright filtered light, complete wetting, rapid drying, and open air circulation form the practical foundation.

LIGHT

Give bright indirect light. Silver xeric types usually accept more sun than soft green mesic types, but acclimate every plant gradually.

WATER

Wet the entire plant by rinsing or soaking, then shake gently and orient it so trapped water can escape.

AIRFLOW

Let the plant dry fully in open air before it returns to a vessel or display. Avoid sealed glass.

WARMTH

Days around 65–90°F and nights above 55°F suit many common types. Avoid heaters, air conditioners, and frost.

DISPLAY

Keep the base unburied and uncovered. Use an open support that does not trap water against the leaves.

UPKEEP

Remove only loose dead basal leaves, inspect the center after watering, and watch for pups after flowering.

THE RELIABLE DEFAULT

Wet every leaf, drain the center, and return the plant to its display only after it has dried.

Light

Window direction is only the beginning. Trees, buildings, season, curtains, and distance all change the light that reaches each frond.

TOO INTENSE

Pale, scorched, or crisp patches after exposure to hot direct sun.

USEFUL RANGE

Bright indirect light supports compact growth and flowering. An east or west window is often a useful starting point.

TOO DIM

Weak stretched growth, fading color, slow drying, or a plant that never approaches bloom.

ADJUST SLOWLY

Move the plant in stages. Leaves formed in shade can scorch after a sudden move into stronger light.

READ THE WHOLE PLANT

Dense silvery trichomes usually signal a type adapted to brighter, drier conditions. Soft green leaves generally call for gentler light and more frequent moisture.



Placement

Choose an open display near bright filtered light. Leave air around every rosette and remove the plants for watering so the mount and leaf bases can dry.

Watering

Rinsing, misting to runoff, and soaking can all work. The right frequency depends on species, light, humidity, heat, and drying speed.



After rinsing or soaking, shake out trapped water and let the plants dry in open air before returning them to display.

01

CHECK BELOW

Notice whether leaves curl more tightly, feel unusually light, or lose their normal flexibility. Compare with the plant's usual dry state.

02

WATER EVENLY

Rinse thoroughly or soak for roughly 20–60 minutes, following the response of the species and room rather than a rigid calendar.

03

LET IT FINISH

Shake gently, angle the opening downward, and provide moving air. Do not return it to enclosed glass until fully dry.

Humidity and temperature

Humidity changes how often a *Tillandsia* needs water, but airflow determines whether the plant dries safely afterward.

WARM ROOMS	Penn State recommends daytime temperatures of 65–90°F (18–32°C) and nights above 55°F (13°C). Protect plants from frost and hot glass.
HUMIDITY	Dry rooms usually require more frequent wetting. Humid rooms require less, but the leaf bases must still dry with moving air.
WINTER	Heating vents can dehydrate leaves while dim winter light slows drying. Move the plant away from forced air and reassess both factors.

DRYING IS PART OF WATERING

Water thoroughly, then give the center and leaf bases enough air to become fully dry.

Display and feeding



An open display supports the plant without burying its base or trapping water between the leaves.

OPEN SUPPORT

Use cork, clean stone, open wire, or a shallow unglazed support. Keep the plant removable for watering and leave the rosette exposed to air.

FERTILIZER

Fertilizer is optional. During active growth, use a bromeliad-compatible fertilizer at a small fraction of label strength and never let concentrate collect in the center.

Air plant anatomy



Numbered markers correspond to the definitions at right.

1 ROSETTE

The spiral arrangement of leaves around the plant's short central stem.

2 LEAF

A simple, usually narrow blade that channels water toward the center and carries absorptive trichomes.

3 TRICHOMES

Tiny absorptive scales that give many species a silver or fuzzy surface.

4 BASE

The crowded center where leaf bases overlap; trapped water here can encourage decay.

5 ANCHORING ROOTS

Roots mainly fasten an epiphytic plant to bark or rock; they are not planted in ordinary soil.

6 INFLORESCENCE

The flower-bearing stalk that rises from the rosette, often with colorful bracts.

Drying and grooming



DRAIN THE CENTER

After watering, shake gently and angle the rosette downward so water does not remain pooled between overlapping leaf bases.

GIVE IT OPEN AIR

Dry the plant in a bright, ventilated place. Penn State advises at least four hours of drying before returning it to a terrarium.

REMOVE ONLY LOOSE TISSUE

Gently pull away a fully dry basal leaf only when it releases easily. Do not strip living tissue or damage a developing pup.

PROTECT THE TRICHOMES

Do not scrub or polish silvery leaves. Their pale surface is functional tissue, not dust.

JUDGE THE CENTER FIRST

Crisp tips point toward dehydration or excess salts. A soft, darkening base suggests trapped moisture and requires faster drying.

Troubleshooting

Check the leaf texture, the base, drying time, and the most recent environmental change before choosing a response.

WHAT YOU SEE	WHAT TO INSPECT	FIRST RESPONSE
Crisp brown tips	Long gaps between wettings, hot dry airflow, hard-water salts, or excess fertilizer.	Restore thorough wetting, move from vents, and pause fertilizer.
Tighter leaf curl	Normal species form versus a recent increase in curl, lighter weight, or loss of flexibility.	Water thoroughly and reassess after the plant has dried.
Soft, dark base	Water trapped in the rosette, slow drying, enclosed glass, or poor airflow.	Remove from the display, dry in open air, and remove only tissue that is clearly dead.
Pale or scorched patches	Hot direct sun, reflected heat, or a sudden move from shade into stronger light.	Move back from the glass and adjust light gradually.
Scale or mealybugs	Leaf bases, the sheltered center, and the underside of leaves.	Separate the plant, confirm the pest, and follow a labeled treatment suitable for bromeliads.

DO NOT STACK REMEDIES

Extra soaking, fertilizer, pruning, pesticide, and a new display applied together can increase stress and erase the evidence. Change one condition at a time unless the base is actively failing.

Mounting and remounting



A mount should support the plant without burying its base, crushing leaves, or preventing removal for watering.

01

CHOOSE

Select a dry, open support such as cork, clean stone, or unglazed ceramic with room around the rosette.

02

POSITION

Rest the base against the support without pushing it into soil, moss, or a deep vessel.

03

SECURE

Use a loose plant-safe tie only if needed. Keep pressure away from the center, new leaves, and any pup.

04

KEEP REMOVABLE

Lift the plant away for watering and return it only after both plant and support are fully dry.

ANCHORING ROOTS ARE NORMAL

Fine roots may grip a mount over time. Leave healthy attached roots in place; they support the epiphytic habit even though most moisture is absorbed through the leaves.

Propagation

Each rosette blooms once, then can produce offsets called pups. Leaving pups attached creates a natural clump; separation is optional.



Pups arise from the parent plant's base. Let them grow substantially before deciding whether to separate them.

01

CHOOSE

Wait until the parent has flowered and an offset is clearly established at the base.

02

GROW

Allow the pup to reach roughly one-third to one-half of the parent's size. Larger pups recover more reliably.

03

SEPARATE

Ease a mature pup sideways from the base, preserving its basal tissue. Do not tear into the center of the parent.

04

RECOVER

Keep the pup in bright filtered light, wet it thoroughly, and let it dry fully after every watering.

LEAVE THE CLUMP IF YOU PREFER

Attached pups are not a problem. A multi-rosette cluster often becomes fuller and more stable than newly separated plants.

Seasonal care

Indoor seasons change light, drying time, humidity, and growth. Adjust watering from observation rather than the calendar alone.

OBSERVE	SPRING	SUMMER	AUTUMN	WINTER
LIGHT	Increase filtered light gradually.	Protect from hot midday sun.	Reassess as daylight shortens.	Use the brightest safe position.
WATER	Resume more frequent full wettings.	Check more often in heat and dry air.	Adjust as rooms cool.	Water less often if drying slows.
FEED	Begin reduced-strength feeding.	Feed periodically during active growth.	Reduce as growth slows.	Do not fertilize.
PROJECTS	Check mounts and developing pups.	Move outdoors only to bright shade.	Inspect before bringing indoors.	Protect from vents and dry heat.

THE USEFUL SCHEDULE

Schedule observations. Let the observations determine the care.

Pet safety

This guide covers many *Tillandsia* species. Treat pet safety as species-specific unless your plant is positively identified and checked in a veterinary toxicology source.

PRACTICAL PRECAUTIONS

Do not assume that the broad store label “air plant” establishes identity or safety. Keep the nursery label when one is provided.

Collect detached leaves, spent flowers, loose ties, wire, and mounting debris promptly.

Prevent access to hanging hardware, glass vessels, or brittle ceramic supports that can cause injury if pulled down.

Keep fertilizers, pesticides, adhesives, and mounting materials secured.

WHAT TO DO

Chewing plant material can still cause stomach upset, and fertilizers or pesticides may be harmful. Contact a veterinarian if a pet develops symptoms or may have swallowed plant-care products.

ASPCA ANIMAL POISON CONTROL

(888) 426-4435

A consultation fee may apply.

Keep the species name and any product labels available when seeking veterinary advice. A genus-level care guide cannot establish the safety of every plant sold under a common name.

Observation log: before a change

Use one page for one care decision. Print extra copies of pages 18-19 whenever you need them.

PLANT NAME OR IDENTIFYING NOTE

DATE

LOCATION AND WINDOW DIRECTION

WHAT I OBSERVED

PLANT STATE BEFORE WATERING

☐ Fully dry ☐ Slightly flexible ☐ More curled ☐ Very light ☐ Not checked

ONE ACTION I TOOK

Observation log: follow-up

Return after enough time has passed to notice a meaningful response. Avoid adding a second change too quickly.

FOLLOW-UP DATE

DAYS SINCE THE CHANGE

WHAT CHANGED NEXT

WHAT STAYED THE SAME

MY NEXT OBSERVATION DATE AND WHY

☐ Continue ☐ Reassess ☐ Seek expert help ☐ No further action

Continue observing

Keep the guide. Reprint the log. Let observation become the routine.