

The Spider Plant Guide

Chlorophytum comosum



How to use this guide

The arch of the leaves. The color of the tips. The weight of the pot. The appearance of new plantlets. Good care begins by noticing what changed before deciding what to do.

THE ONE-CHANGE RULE

Change one condition.
Record what happened. Give
the plant time to answer.

Contents

PLANT IDENTITY

BOTANICAL *Chlorophytum
comosum*

FAMILY Asparagaceae

ORIGIN Tropical and
southern Africa

HABIT Clump-forming,
rhizomatous perennial

SECTION	PAGE
Plant profile	04
Care at a glance	05
Light	06
Placement	07
Watering	08
Humidity and temperature	09
Soil and fertilizer	10
Spider plant anatomy	11

SECTION	PAGE
Leaf care and cleanup	12
Troubleshooting	13
Repotting	14
Propagation	15
Seasonal care	16
Pet safety	17
Observation log	18

HOW TO USE THIS GUIDE

Start with the condition you can observe. Avoid making several changes at once.
The printable log at the end is part of the care system, not an afterthought.



Plant profile

Spider plant is a clump-forming perennial with arching strap-shaped leaves. Slender flowering stems extend beyond the foliage and can carry small white flowers followed by young plantlets.

Care at a glance

Bright, lightly shaded, evenly tended, and observed. Spider plant is adaptable, but its leaf tips and growth still record the care it receives.

LIGHT

Medium to bright indirect light supports dense growth and plantlets. Protect the leaves from harsh direct sun.

WATER

Allow the mix to dry slightly, then water thoroughly and let excess drain. Do not keep the roots continuously wet.

HUMIDITY

Average household humidity usually works. Dry air can contribute to brown tips, but saturated soil is the greater risk.

WARMTH

Cool to average indoor temperatures suit it. Protect the plant from frost, cold glass, and drying heat.

SOIL

Use a general-purpose, well-drained indoor mix in a container with drainage holes.

UPKEEP

Trim fully brown tips or damaged leaves cleanly, and remove spent flowering stems when they decline.

THE RELIABLE DEFAULT

Let the surface dry slightly, then water thoroughly and empty the saucer or outer pot.

Light

Window direction is only the beginning. Trees, buildings, season, curtains, and distance all change the light that reaches a leaf.

TOO INTENSE

Pale, bleached, or scorched areas after exposure to strong direct sun.

USEFUL RANGE

Medium to bright filtered daylight that keeps variegation clear without scorching the leaves.

TOO DIM

Slower growth, sparse plantlets, weaker variegation, or a clump leaning toward the window.

ADJUST SLOWLY

Move the plant in stages. Sudden changes can burn or stress established leaves.

WATCH THE WHITE STRIPES

Variegated leaves make light stress easier to see. Protect pale tissue from strong sun and move the plant gradually when changing locations.



Placement

Give the arching leaves and runners room to hang without rubbing against glass, heaters, or foot traffic. Filter strong sun and adjust the location gradually.

Watering

The calendar should remind you to inspect the pot, not instruct you to pour.



Water the whole root zone, then let the container drain completely.

01

CHECK THE MIX

Feel below the surface and lift the pot. If the mix still feels moist and the pot feels heavy, wait.

02

WATER EVENLY

When the surface has dried slightly, wet the mix across the pot instead of pouring in one spot.

03

LET IT FINISH

Drain completely and empty any saucer or decorative outer pot.

Humidity and temperature

Spider plant handles ordinary indoor air, though very dry conditions can contribute to crisp leaf tips.

GOOD RANGE	Cool to average indoor temperatures are suitable. Keep the plant above 45°F (7°C) and away from cold drafts or hot vents.
HUMIDITY	Average household humidity is usually adequate. A humidifier can help in persistently dry rooms, but do not compensate by keeping the mix wet.
WINTER	Shorter days slow growth and water use. Check the mix rather than maintaining the same summer watering interval.

SEPARATE AIR FROM SOIL

Humidity may affect the tips, but extra water in the pot does not raise room humidity safely.

Soil and fertilizer



The potting mix supports the crown while leaving a small gap below the rim.


POTTING MIX

Choose a general-purpose potting soil or soilless indoor medium. Avoid dense garden soil and containers without drainage.

FERTILIZER

Spider plant needs modest feeding. Follow the product label, or use a diluted fertilizer occasionally during active growth; excess fertilizer can reduce plantlets and contribute to brown tips.

Spider plant anatomy



Numbered markers correspond to the definitions at right.

1 LEAF BLADE

The arching, strap-shaped photosynthetic surface.

2 CENTRAL STRIPE

The pale lengthwise band typical of the cultivar shown.

3 BASAL ROSETTE

The crown where arching leaves emerge in a dense cluster.

4 FLOWERING STEM

A slender arching stem that bears small flowers and later plantlets.

5 PLANTLET

A young rosette that develops along the flowering stem and can form roots.

6 FLESHY ROOTS

Thickened pale roots that store water alongside finer absorbing roots.

Leaf tips and cleanup



READ THE PATTERN

One brown tip may be old damage. Many new brown tips suggest a repeated issue such as dryness, salts, or water quality.

TRIM CONSERVATIVELY

Use clean scissors to remove only dead brown tissue, following the natural leaf shape without cutting into green tissue.

REMOVE FAILED LEAVES

Cut a fully yellow or badly damaged leaf near the base. Do not shorten healthy leaves simply to make the clump even.

CLEAN WITHOUT COATING

Rinse dust gently or wipe leaves with plain water. Skip commercial leaf-shine products.

A TRIM IS COSMETIC

Removing the brown portion improves appearance but does not correct the cause. Review moisture, fertilizer, room dryness, and water quality.

Troubleshooting

Check the pattern, the root zone, and the recent change before reaching for a remedy.

WHAT YOU SEE	WHAT TO INSPECT	FIRST RESPONSE
Brown leaf tips	Dry air, prolonged dryness, accumulated fertilizer salts, or sensitivity to tap-water chemicals.	Review the pattern and recent care. Correct one likely cause before trimming dead tissue.
Pale or scorched leaves	Strong direct sun, especially after a sudden move to a brighter window.	Move back from the glass or filter the light, then allow new growth to show the response.
Yellowing and soft growth	Persistently wet mix, blocked drainage, cold conditions, or declining roots.	Pause watering and inspect the pot and roots before adding fertilizer or more moisture.
Few plantlets	Low light, heavy feeding, an immature plant, or a recent change in growing conditions.	Provide steady medium-to-bright indirect light and avoid pushing growth with extra fertilizer.
Scale or cottony pests	Leaf bases, undersides, and runners for scale insects or mealybug clusters.	Separate the plant, confirm the pest, and follow a labeled treatment.

DO NOT STACK REMEDIES

Extra water, fertilizer, repotting, homemade pesticide, and a new location applied together can increase stress and erase the evidence. Change one condition at a time unless roots are actively failing.

Repotting



Fleshy roots can fill a container quickly even while the foliage looks comfortable.

01

PREPARE

Choose a draining pot one size larger and prepare fresh mix.

02

INSPECT

Look before teasing. Healthy storage roots are firm and pale, not soft or foul-smelling.

03

SET

Keep the crown at its original depth and spread only roots that loosen without force.

04

SETTLE

Backfill gently, water once, drain, and return to stable light.

WHEN TO ACT

A slightly snug pot can support plantlet production, but severely crowded roots, distorted containers, or mix that no longer accepts water are signals to act. Move up one pot size or divide the clump. Active rot or blocked drainage should not wait for spring.

Propagation

Plantlets are ready-made young rosettes. Root one while it remains attached to the mother plant, then cut the runner after roots establish.



Rooting an attached plantlet reduces stress while the young plant establishes.

01

CHOOSE

Select a healthy plantlet that has begun to form small aerial roots.

02

SET

Place its base on moist soilless medium in a small draining pot beside the mother plant.

03

ANCHOR

Hold the plantlet gently against the surface with bent wire if needed. Keep the crown above the mix.

04

SEPARATE

After roots form, cut the runner and care for the young plant on its own.

Seasonal care

Indoor seasons are shaped by daylight, heating, cooling, and windows. The plant's pace remains the primary signal.

OBSERVE	SPRING	SUMMER	AUTUMN	WINTER
LIGHT	Increase light gradually.	Filter strong sun.	Watch shorter days.	Use the brightest safe place.
WATER	Growth and drying may accelerate.	Check moisture more often.	Intervals may widen.	Confirm dryness first.
FEED	Begin with growth.	Continue at label rate.	Reduce as growth slows.	Pause if inactive.
PROJECTS	Repot, divide, or root plantlets.	Watch runners and brown tips.	Remove tired stems and inspect pests.	Stabilize warmth.

THE USEFUL SCHEDULE

Schedule observations. Let the observations determine the care.

Pet safety

The ASPCA lists spider plant as non-toxic to cats and dogs.

PRACTICAL PRECAUTIONS

Place hanging baskets where pets cannot pull them down.

Redirect repeated chewing to protect both pet and plant.

Collect fallen plantlets and trimmed leaves promptly.

Keep fertilizers, pesticides, and potting materials secured.

WHAT TO DO

Non-toxic does not mean every exposure is harmless. Contact your veterinarian if a pet develops persistent symptoms, eats potting materials, or may have contacted fertilizer or pesticide.

ASPCA ANIMAL POISON CONTROL

(888) 426-4435

A consultation fee may apply.

The non-toxic classification applies to the plant itself. Products used on the plant and materials in the container require their own safety precautions.

Observation log: before a change

Use one page for one care decision. Print extra copies of pages 18-19 whenever you need them.

PLANT NAME OR IDENTIFYING NOTE

DATE

LOCATION AND WINDOW DIRECTION

WHAT I OBSERVED

POTTING MIX BEFORE WATERING

☐ Wet
 ☐ Moist
 ☐ Dry at surface
 ☐ Dry deeper
 ☐ Not checked

ONE ACTION I TOOK

Observation log: follow-up

Return after enough time has passed to notice a meaningful response. Avoid adding a second change too quickly.

FOLLOW-UP DATE

DAYS SINCE THE CHANGE


WHAT CHANGED NEXT

WHAT STAYED THE SAME

MY NEXT OBSERVATION DATE AND WHY

☐ Continue ☐ Reassess ☐ Seek expert help ☐ No further action

Continue observing



Keep the guide. Reprint the log. Let observation become the routine.